

# Package ‘ICSS’

January 20, 2025

**Type** Package

**Title** ICSS Algorithm by Inclan/Tiao (1994)

**Version** 1.1

**Date** 2021-04-22

**Maintainer** Siegfried Köstlmeier <[siegfried.koestlmeier@gmail.com](mailto:siegfried.koestlmeier@gmail.com)>

**Description** The Iterative Cumulative Sum of Squares (ICSS) algorithm by Inclan/Tiao (1994) <<https://www.jstor.org/stable/2290916>> detects multiple change points, i.e. structural break points, in the variance of a sequence of independent observations. For series of moderate size (i.e. 200 observations and beyond), the ICSS algorithm offers results comparable to those obtained by a Bayesian approach or by likelihood ratio tests, without the heavy computational burden required by these approaches.

**License** GPL-2

**Depends** R (>= 3.5.0)

**Imports** rstack

**Suggests** testthat

**Encoding** UTF-8

**LazyData** true

**NeedsCompilation** no

**Author** Siegfried Köstlmeier [aut, cre, cph]  
(<<https://orcid.org/0000-0002-7221-6981>>)

**Repository** CRAN

**Date/Publication** 2021-04-22 14:00:19 UTC

## Contents

data . . . . .	2
ICSS . . . . .	2

## Index

4

data

*Sample data for Inclan/Tiao (1994)***Description**

Generated random data (n=700) with following the scheme in Inclan/Tiao (1994):

- [1;390]Mean: 0; Variance: 1.000
- [391;517]Mean: 0; Variance: 0.365
- [518;700]Mean: 0; Variance: 1.033

**Usage**

```
data(data)
```

**Examples**

```
## load data
data(data)

## calculate the variance until the first breakpoint.
data_var <- var(data[1:390])
```

ICSS

*Iterative Cumulative Sum of Squares (ICSS)***Description**

ICSS implements the Iterative Cumulative Sum of Squares (ICSS) algorithm by Inclan/Tiao (1994). The test detects structural breakpoints in the variance of time series data.

**Usage**

```
ICSS(data, demean = FALSE)
```

**Arguments**

data	A numerical vector
demean	An object of class " <a href="#">logical</a> ": If demean is TRUE, all data will get demeaned prior to the ICSS algorithm.

**Value**

ICSS returns a numerical vector containing the location of structural breakpoints or NA if none breakpoints are found.

## References

Inclan, C., & Tiao, G. C. (1994): Use of cumulative sums of squares for retrospective detection of changes of variance. *Journal of the American Statistical Association*, 89(427), 913-923. <https://www.jstor.org/stable/2290916>.

## Examples

```
## load demo data
data(data)
breakpoints <- ICSS(data)
```

# Index

[data, 2](#)

[ICSS, 2](#)

[logical, 2](#)